

AD-A115 280

FLORIDA STATE UNIV TALLAHASSEE DEPT OF STATISTICS

F/G 12/1

THE RELIABILITY OF K OUT OF N SYSTEMS.(U)

MAR 82 P J BOLAND, F PROSCHAN

F49620-82-K-0007

UNCLASSIFIED

FSU-STATISTICS-M610

AFOSR-TR-82-0453

NL

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111

111


111

END

DATE

7 82

DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER <b>AFOSR-TR- 82-0453</b>	2. GOVT ACCESSION NO. <b>AD-A115280</b>	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) <b>THE RELIABILITY OF k OUT OF n SYSTEMS</b>		5. TYPE OF REPORT & PERIOD COVERED <b>TECHNICAL</b>
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) <b>Philip J. Boland and Frank Proschan*</b>		8. CONTRACT OR GRANT NUMBER(s) <b>F49620-82-K-0007</b>
9. PERFORMING ORGANIZATION NAME AND ADDRESS <b>*Department of Statistics, Florida State Univ, Tallahassee FL 32306; and Department of Mathema- tics, University College Dublin, Belfield, Dublin 4, Ireland</b>		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS <b>PE61102F; 2304/A5</b>
11. CONTROLLING OFFICE NAME AND ADDRESS <b>Directorate of Mathematical &amp; Information Sciences Air Force Office of Scientific Research Bolling AFB DC 20332</b>		12. REPORT DATE <b>MAR 82</b>
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES <b>11</b>
		15. SECURITY CLASS. (of this report) <b>UNCLASSIFIED</b>
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) <b>Approved for public release; distribution unlimited.</b>		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) <b>k out of n systems, majorization, Schur-convexity and Schur-concavity, independent Bernoulli trials.</b> <b>P-sub1 P-subn h-subk</b>		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) <b>A system with n independent components which function if and only if at least k of the components function is a k out of n system. Parallel systems are 1 out of n systems and series systems are n out of n systems. If <math>p = (p_1, \dots, p_n)</math> is the vector of component reliabilities for the n components, then <math>h_k(p)</math> is the reliabi- lity function of the system. It is shown that <math>h_k(p)</math> is Schur-convex in</b>		

AD A115280

DTIC FILE COPY

(Continued)

DD FORM 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED  
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

ITEM #20, CONTINUED:

$\left[ \frac{k-1}{n-1}, 1 \right]^n$ . More particularly if  $\Pi$  is an  $n \times n$  doubly stochastic matrix, then  $h_k(p) \geq (\leq) h_k(p\Pi)$  whenever  $p \in \left[ \frac{k-1}{n-1}, 1 \right]^n \left( \left[ 0, \frac{k-1}{n-1} \right]^n \right)$ . This theorem is compared with a result on Schur-convexity and -concavity by Gleser [2] which in turn extends work of Hoeffding [4].



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Avail and/or	
Dist	Special
A	

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

**AFOSR-TR- 82 - 0 4 5 3**

**The Reliability of  $k$  out of  $n$  Systems**

by

**Philip J. Boland and Frank Proschan<sup>1</sup>**

**University College Dublin  
Department of Mathematics  
Belfield, Dublin 4, Ireland**

and

**The Florida State University  
Department of Statistics  
Tallahassee, Florida 32306**

*STATISTICS*  
**FSU Report M610  
AFOSR Technical Report 82-136**

<sup>1</sup>**Research supported by Air Force Office of Scientific Research,  
U.S.A.F., under Contract No. F49620-82-K-0007.**

**AMS Subject Classification Numbers: Primary 60C05, 62N05;  
Secondary 62E15, 26B25.**

**Key Words and Phrases:  $k$  out of  $n$  systems, majorization,  
Schur-convexity and Schur-concavity, independent Bernoulli  
trials.**

**Approved for public release;  
distribution unlimited.**

**82 00 02 056**

## The Reliability of k out of n Systems

by

Philip J. Boland and Frank Proschan

A system with  $n$  independent components which functions if and only if at least  $k$  of the components function is a  $k$  out of  $n$  system. Parallel systems are 1 out of  $n$  systems and series systems are  $n$  out of  $n$  systems. If  $\underline{p} = (p_1, \dots, p_n)$  is the vector of component reliabilities for the  $n$  components, then  $h_k(\underline{p})$  is the reliability function of the system. It is shown that  $h_k(\underline{p})$  is Schur-convex in  $\left[\frac{k-1}{n-1}, 1\right]^n$  and Schur-concave in  $\left[0, \frac{k-1}{n-1}\right]^n$ . More particularly if  $\Pi$  is an  $n \times n$  doubly stochastic matrix, then  $h_k(\underline{p}) \geq (\leq) h_k(\underline{p}\Pi)$  whenever  $\underline{p} \in \left[\frac{k-1}{n-1}, 1\right]^n$   $\left(\left[0, \frac{k-1}{n-1}\right]^n\right)$ . This Theorem is compared with a result on Schur-convexity and -concavity by Gleser [2] which in turn extends work of Hoeffding [4].

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)  
NOTICE OF TRANSMITTAL TO DTIC  
This technical report has been reviewed and is  
approved for public release IAW AFR 190-12.  
Distribution is unlimited.  
MATTHEW J. KERPER  
Chief, Technical Information Division

1. Introduction and Summary. A  $k$  out of  $n$  system is a system with  $n$  components which functions if and only if  $k$  or more of the components function. Herein we assume that the  $n$  components of the system function independently. If  $\underline{p} = (p_1, \dots, p_n)$  is the vector of component reliabilities (functioning probabilities) and  $\underline{\epsilon} = (\epsilon_1, \dots, \epsilon_n)$  represents any vector with components equal to zeroes or ones, then

$$(1.1) \quad h_k(\underline{p}) = h_k(p_1, \dots, p_n) = \sum_{\substack{\underline{\epsilon}, \epsilon_1, \dots, \epsilon_n \\ \epsilon_1 + \dots + \epsilon_n \geq k}} p_1^{\epsilon_1} \dots p_n^{\epsilon_n} (1-p_1)^{1-\epsilon_1} \dots (1-p_n)^{1-\epsilon_n}$$

is the probability that  $k$  or more of the components function. This function  $h_k: [0, 1]^n \rightarrow [0, 1]$  is called the reliability function for a  $k$  out of  $n$  system with independent components. A one out of  $n$  system is a parallel system, an  $n - 1$  out of  $n$  system is a 'fail-safe' system (see Barlow-Proschan [1]), and an  $n$  out of  $n$  system is a series system. For these systems it is easy to see that

$$h_1(\underline{p}) = 1 - \prod_{i=1}^n (1 - p_i),$$

$$h_{n-1}(\underline{p}) = \prod_{i=1}^n p_i + \prod_{i=1}^n ((1 - p_i) \prod_{j=1}^n p_j)$$

and

$$h_n(\underline{p}) = \prod_{i=1}^n p_i.$$

Note that if  $S$  is the number of successes in  $n$  independent Bernoulli

trials, where for  $i = 1, \dots, n$ ,  $p_i$  is the probability of success on the  $i$ th trial, then  $P(S = k - 1) = h_k(p) = h_k(p_1, \dots, p_n)$ . Hence the results in this paper on  $k$  out of  $n$  systems have interpretations in terms of the number of successes in Bernoulli trials.

If  $p = (p_1, \dots, p_n)$ , then let  $\bar{p} = (\sum_{i=1}^n p_i/n, \dots, \sum_{i=1}^n p_i/n)$ . Hoeffding ([4], 1956) showed that

$$h_k(p_1, \dots, p_n) \geq h_k(\bar{p}, \dots, \bar{p}) \quad \text{if} \quad \sum_{i=1}^n p_i \geq k$$

and

$$h_k(p_1, \dots, p_n) \leq h_k(\bar{p}, \dots, \bar{p}) \quad \text{if} \quad \sum_{i=1}^n p_i \leq k - 1.$$

Gleser [2], using the theory of majorization and Schur functions (see Marshall and Olkin [5], Theorem 12. K.1), extended Hoeffding's result and showed that  $h_k(p)$  is Schur-convex in the region where  $\sum_{i=1}^n p_i \geq k + 1$  and Schur-concave in the region where  $\sum_{i=1}^n p_i \leq k - 2$ . More particularly Gleser showed that if  $p = (p_1, \dots, p_n) \in [0, 1]^n$  and  $\Pi$  is a doubly stochastic matrix, then

$$h_k(p) \geq h_k(p\Pi) \quad \text{whenever} \quad \sum_{i=1}^n p_i \geq k + 1$$

and

$$h_k(p) \leq h_k(p\Pi) \quad \text{whenever} \quad \sum_{i=1}^n p_i \leq k - 2.$$

This result allows one to make more general comparisons than one could with Hoeffding's result. The major result of the present paper enables one to extend the regions of Schur-convexity and -concavity



of the function  $h_k(p)$ . The result is as follows:

**Theorem 1.1.**  $h_k(p)$  is Schur-concave in the region  $\left[0, \frac{k-1}{n-1}\right]^n$  and Schur-convex in the region  $\left[\frac{k-1}{n-1}, 1\right]^n$ .

2. Majorization, Schur-convexity, and Schur-concavity. A vector

$\underline{x} = (x_1, \dots, x_n)$  is said to majorize a vector  $\underline{y} = (y_1, \dots, y_n)$  ( $\underline{x} \succ \underline{y}$ ) if  $x_{[1]} \geq y_{[1]}$ ,  $x_{[1]} + x_{[2]} \geq y_{[1]} + y_{[2]}$ , ...,  $\sum_{i=1}^{n-1} x_{[i]} \geq \sum_{i=1}^{n-1} y_{[i]}$ , and  $\sum_{i=1}^n x_{[i]} = \sum_{i=1}^n y_{[i]}$ , where the  $x_{[i]}$ 's and  $y_{[i]}$ 's are components of  $\underline{x}$  and  $\underline{y}$  respectively arranged in descending order. The following lemma characterizes majorization and is due to Hardy, Littlewood and Pólya ([3], 1934) (see also Marshall and Olkin [5], Theorem 2.8.2).

**Lemma 2.1.** The vector  $\underline{x}$  majorizes the vector  $\underline{y}$  if and only if there exists an  $n \times n$  doubly stochastic matrix  $\Pi$  such that  $\underline{y} = \underline{x}\Pi$ .

A real valued function  $h$  defined on a set  $A \subset \mathbb{R}^n$  is Schur-convex (Schur-concave) if  $h(\underline{x}) \geq (\leq) h(\underline{y})$  whenever  $\underline{x} \succ \underline{y}$  and  $\underline{x}, \underline{y} \in A$ . Now assume that  $A \subset \mathbb{R}^n$  is a permutation symmetric convex set with nonempty interior. If  $h$  is continuously differentiable on the interior  $A^\circ$  of  $A$  and continuous on  $A$  then  $h$  is Schur-convex (-concave) on  $A$  if and only if for all  $i \neq j$  and  $\underline{x} \in A^\circ$ ,

$$(2.1) \quad (x_i - x_j) \left( \frac{\partial h}{\partial x_i}(\underline{x}) - \frac{\partial h}{\partial x_j}(\underline{x}) \right) \geq (\leq) 0.$$

This characterization of Schur-convexity (-concavity) is known as the Schur-Ostrowski condition (see Marshall and Olkin [5]).

In investigating regions of Schur-convexity of the reliability function  $h_k: [0, 1]^n \rightarrow [0, 1]$ , we use the following notation. If  $\underline{x} \in [0, 1]^m$  and  $k$  is any integer then  $h_k(\underline{x})$  ( $h_k^*(\underline{x})$ ) will denote the probability that  $k$  or more (exactly  $k$ ) independent components with respective probabilities  $x_1, \dots, x_m$  function. Note for example that with this notation  $h_{-1}(\underline{x}) = 1$  and  $h_{-1}^*(\underline{x}) = 0$ . We assume here that  $\underline{p} = (p_1, \dots, p_n) \in [0, 1]^n$  and let  $\underline{p}^i$  ( $\underline{p}^{ij}$ ) be the vector in  $[0, 1]^{n-1}$  ( $[0, 1]^{n-2}$ ) obtained from  $\underline{p}$  by deleting its  $i$ th coordinate ( $i$ th and  $j$ th coordinates).

Lemma 2.2. For  $1 \leq k < n$ ,  $\underline{p} \in (0, 1)^n$  and  $n > 2$ ,

$$\left( \frac{\partial h_k}{\partial p_i}(\underline{p}) - \frac{\partial h_k}{\partial p_j}(\underline{p}) \right) (p_i - p_j) = -(p_i - p_j)^2 [h_{k-2}^*(\underline{p}^{ij}) - h_{k-1}^*(\underline{p}^{ij})]$$

for all  $i, j$ ,  $i \neq j$ .

Proof. For any index  $i$ ,  $1 \leq i \leq n$ , we have that

$$h_k(\underline{p}) = p_i h_{k-1}(\underline{p}^i) + (1 - p_i) h_k(\underline{p}^i).$$

$$\text{Hence for } j \neq i, \frac{\partial h_k}{\partial p_i}(\underline{p}) = h_{k-1}(\underline{p}^i) - h_k(\underline{p}^i)$$

$$= p_j h_{k-2}(\underline{p}^{ij}) + (1 - p_j) h_{k-1}(\underline{p}^{ij})$$

$$- (p_j h_{k-1}(\underline{p}^{ij}) + (1 - p_j) h_k(\underline{p}^{ij})).$$

$$\begin{aligned}
 \text{Therefore } & \left( \frac{\partial h_k}{\partial p_i}(p) - \frac{\partial h_k}{\partial p_j}(p) \right) (p_i - p_j) \\
 &= -(p_i - p_j)^2 [(h_{k-2}^{ij}(p) - h_{k-1}^{ij}(p)) - (h_{k-1}^{ij}(p) - h_k^{ij}(p))] \\
 &= -(p_i - p_j)^2 [h_{k-2}^{*ij}(p) - h_{k-1}^{*ij}(p)].
 \end{aligned}$$

We see then by the Schur-Ostrowski condition (2.1) that  $h_k(p)$  is Schur-convex (-concave) in regions where for all  $i, j, i \neq j$ ,  $h_{k-2}^{*ij}(p) \leq (\geq) h_{k-1}^{*ij}(p)$ . In particular note that if  $k = 1$  (respectively  $n$ ), i.e. for a parallel (series) system,  $h_{k-2}^{*ij}(p) = 0$  ( $h_{k-1}^{*ij}(p) = 0$ ), and hence  $h_1(p)$  ( $h_n(p)$ ) is Schur-convex (-concave) in  $[0, 1]^n$ .

**Lemma 2.3.** Let  $n > 2$  and  $k$  be such that  $2 \leq k \leq n$ . Then for any  $i \neq j$ ,

$$h_{k-2}^{*ij}(p) \leq (\geq) h_{k-1}^{*ij}(p)$$

whenever  $p_l \geq (\leq) \frac{k-1}{n-1}$  for all  $l \neq i, j$ .

**Proof.** Due to the symmetry of the situation, we need only show that

$$h_{k-2}^{*}(p^{12}) \leq h_{k-1}^{*}(p^{12}) \text{ whenever } p_l \geq \frac{k-1}{n-1} \text{ for } l = 3, \dots, n.$$

In this proof,  $\underline{\epsilon} = (\epsilon_3, \dots, \epsilon_n)$  and  $\underline{\epsilon'} = (\epsilon'_3, \dots, \epsilon'_n)$  will denote  $n-2$  dimensional vectors whose components are zeroes or ones such that  $\epsilon_3 + \dots + \epsilon_n = k-2$  and  $\epsilon'_3 + \dots + \epsilon'_n = k-1$ . Now

$$(2.2a) \quad h_{k-2}^{*}(p^{12}) = \sum_{\underline{\epsilon}} p_3^{\epsilon_3} \dots p_n^{\epsilon_n} (1-p_3)^{1-\epsilon_3} \dots (1-p_n)^{1-\epsilon_n}$$

and

$$(2.2b) \quad h_{k-1}^* (p^{12}) = \sum_{\underline{\epsilon}} p_3^{\epsilon_3} \dots p_n^{\epsilon_n} (1-p_3)^{\epsilon'_3} \dots (1-p_n)^{\epsilon'_n}.$$

Consider a term in the expansion of (2.2a), say for simplicity of notation, the term  $p_3 \dots p_k (1-p_{k+1}) \dots (1-p_n)$ . Then

$$\begin{aligned} & p_3 \dots p_k (1-p_{k+1}) \dots (1-p_n) = \\ & \frac{1}{n-k} \underbrace{[p_3 \dots p_k (1-p_{k+1}) \dots (1-p_n) + \dots + p_3 \dots p_k (1-p_{k+1}) \dots (1-p_n)]}_{n-k \text{ times}} \\ & \leq \frac{1}{n-k} \left[ p_3 \dots p_k \left( \frac{n-k}{n-1} \right) (1-p_{k+2}) \dots (1-p_n) + \dots + p_3 \dots p_k (1-p_{k+1}) \right. \\ & \quad \left. \dots (1-p_{n-1}) \left( \frac{n-k}{n-1} \right) \right] \\ & \quad (\text{since } 1-p_\ell \leq 1 - \frac{k-1}{n-1} = \frac{n-k}{n-1} \text{ for all } \ell = 3, \dots, n) \\ & = \frac{1}{k-1} \left[ p_3 \dots p_k \left( \frac{k-1}{n-1} \right) (1-p_{k+2}) \dots (1-p_n) + \dots + p_3 \dots p_k (1-p_{n+1}) \dots (1-p_{n-1}) \frac{k-1}{n-1} \right] \\ & \leq \frac{1}{k-1} [p_3 \dots p_k p_{k+1} (1-p_{k+2}) \dots (1-p_n) + \dots + p_3 \dots p_k (1-p_{k+1}) \dots (1-p_{n-1}) p_n]. \end{aligned}$$

Hence similarly for a general term in the expansion of (2.2a), we have

$$\begin{aligned} & p_3^{\epsilon_3} \dots p_n^{\epsilon_n} (1-p_3)^{1-\epsilon_3} \dots (1-p_n)^{1-\epsilon_n} \\ & \leq \frac{1}{k-1} \sum_{\substack{\underline{\epsilon'}, \epsilon'_\ell \geq \epsilon_\ell, \ell=3, \dots, n}} p_3^{\epsilon'_3} \dots p_n^{\epsilon'_n} (1-p_3)^{1-\epsilon'_3} \dots (1-p_n)^{1-\epsilon'_n}. \end{aligned}$$

Therefore

$$\begin{aligned}
 h_{k-2}^*(p^{12}) &= \sum_{\underline{\epsilon}} p_3^{\epsilon_3} \dots p_n^{\epsilon_n} (1-p_3)^{1-\epsilon_3} \dots (1-p_n)^{1-\epsilon_n} \\
 &\leq \sum_{\underline{\epsilon}} \frac{1}{k-1} \sum_{\substack{\underline{\epsilon}', \epsilon'_l \geq \epsilon_l, \\ l=3, \dots, n}} p_3^{\epsilon'_3} \dots p_n^{\epsilon'_n} (1-p_3)^{1-\epsilon'_3} \dots (1-p_n)^{1-\epsilon'_n} \\
 &\leq \sum_{\underline{\epsilon}} \frac{1}{k-1} (k-1) p_3^{\epsilon_3} \dots p_n^{\epsilon_n} (1-p_3)^{1-\epsilon_3} \dots (1-p_n)^{1-\epsilon_n} \\
 &\quad \text{(since for each } \underline{\epsilon} \text{ there exist } k-1 \text{ distinct} \\
 &\quad \underline{\epsilon}' \text{ where } \epsilon'_l \geq \epsilon_l \text{ for } l = 3, \dots, n). \\
 &= h_{k-1}^*(p^{12}). \quad \square
 \end{aligned}$$

3. Proof of Theorem 1.1. Using the Schur-Ostrowski condition (2.1) it is easy to verify Theorem 1.1 when  $n = 2$ . For  $n > 2$ , we have already noted that when  $k = 1$  (parallel system),  $h_k$  is Schur-convex in  $[0, 1]^n$ . For  $n > 2$  and  $k \geq 2$ , it follows from Lemmas 2.2 and 2.3 that  $h_k$  satisfies the Schur-Ostrowski condition for Schur-convexity (-concavity) on  $\left[\frac{k-1}{n-1}, 1\right]^n \left[0, \frac{k-1}{n-1}\right]^n$ .

Remark 3.1. Gleser's result [2] shows that  $h_k(p)$  is Schur-convex (-concave) on the set  $\{p: p_1 + \dots + p_n \geq k + 1\}$  ( $\{p: p_1 + \dots + p_n \leq k - 2\}$ ). This result and Theorem 1.1 enable one to make various comparisons of system reliability, and neither result encompasses the other.

For example let us consider a 3 out of 4 system. Then Theorem 1.1 implies that a system with component reliabilities (1.0, .9, .8, .7) is superior (has higher reliability) than a system with component reliabilities (.95, .95, .75, .75) which in turn is superior to one with component reliabilities (.85, .85, .85, .85). On the other hand Theorem 1.1 also implies that a system with component reliabilities (.6, .5, .3, .2) is inferior to one with component reliabilities (.6, .4, .4, .2) which in turn is inferior to one with component reliabilities (.4, .4, .4, .4). These comparisons are not implied by Gleser's result.

Remark 3.2. Let  $S$  be the number of successes in  $n$  independent Bernoulli trials where  $p_i$  is the probability of success on the  $i$ th trial. Then by Theorem 1.1,  $P(S \geq k)$  is Schur convex (concave) in  $\left[\frac{k-1}{n-1}, 1\right]^n \left([0, \frac{k-1}{n-1}]^n\right)$ . Suppose now that  $1 \leq k < k' \leq n$ . Then it follows by the above that  $P(k' > S \geq k)$  is Schur-convex in  $\left[\frac{k-1}{n-1}, \frac{k'-1}{n-1}\right]^n$ .

References.

- [1] Barlow, R.E. and Proschan, F., (1965). Mathematical Theory of Reliability, John Wiley and Sons, New York.
- [2] Gleser, L. (1975). On the distribution of the number of successes in independent trials. Ann. Probab. 3, 182-188.
- [3] Hardy, G.H., Littlewood, J.E., and Pólya, G. (1959). Inequalities, 2nd ed. Cambridge Univ. Press.
- [4] Hoeffding, W. (1956). On the distribution of the number of successes in independent trials. Ann. Math. Statist. 27 713-721.
- [5] Marshall, A.W. and Olkin, I., (1979). Inequalities: Theory of Majorization and its Applications. Academic Press.

DATE  
ILMEI  
—8